

Childhood Immunisation Programme

When to immunise	Diseases protected against	Vaccine given
2 months old	Diphtheria, tetanus, pertussis (whooping cough), polio and Haemophilus influenzae type b (Hib) Pneumococcal disease Rotavirus	DTaP/IPV/Hib and Pneumococcal conjugate vaccine (PCV) Rotarix (Oral)
3 months old	Diphtheria, tetanus, pertussis, polio and Haemophilus influenzae type b (Hib) Meningococcal group C disease (MenC) Rotavirus	DTaP/IPV/Hib and MenC Rotarix (Oral)
4 months old	Diphtheria, tetanus, pertussis, polio and Haemophilus influenzae type b (Hib) Meningococcal group C disease (MenC) Pneumococcal disease	DTaP/IPV/Hib, MenC and PCV
Between 12 and 13 months - within a month of the first birthday	Hib MenC Pneumococcal disease Measles, mumps and rubella (German measles)	Hib/MenC, PCV and MMR
2 years	Flu Vaccine	Fluenz
3 years 4 months to 5 years	Diphtheria, tetanus, pertussis and polio Measles, mumps and rubella	DTaP/IPV or dTaP/IPV and MMR
Girls aged 12 to 13 years	Cervical cancer caused by human papillomavirus types 16 and 18	Gardasil 3 doses
13 to 15 years	Meningitis C	MenC
13 to 18 years old	Tetanus, diphtheria and polio	Td/IPV

Please note that we also offer the HPV vaccination as part of our schools vaccination programme.